Clinical guideline: management of seizures in the emergency room

José Visoso-Franco¹, Liliana Romero-Ocampo², José A. Santos-Zambrano³, Alberto Serrano-González⁴ and Elvira Castro-Martínez⁵

¹Hospital Regional de Especialidades ISSSTE, León, Guanajuato; ²Hospital Central Norte PEMEX y Hospital General de La Raza CMN, Especialidades IMSS; ³Instituto Nacional de Neurología y Neurocirugía; ⁴Hospital Pediátrico Legaria; ⁵Hospital General "Dr. Manuel Gea González" e Instituto Nacional de Neurología y Neurocirugía. Mexico City, Mexico.

Abstract

This clinical guideline on epilepsy contains levels of evidence and recommendations based on the scientific method. Its primary function is to provide emergency medicine physicians a clear diagnostic approach when faced with a pediatric or adult patient with epileptic seizures (ES) or epilepsy. The objective is to unify criteria that will guarantee integral health care, based on adequate decision-making, benefiting the patient through the individualized analysis of proper anamnesis, physical examination, precise indication of laboratory, and image diagnostic tests that yield the pertinent clinical and pharmacological treatment for opportune interventions, avoiding complications, and whenever possible, the recurrence of the ES.

Key words: Seizures. Epilepsy. Treatment in the emergency room.

Introduction

Epilepsy is one of the diseases that affect the quality of life of patients the most, due to its neurological, psychological, and social implications. Epileptic seizures (ES) are a common cause of admittance to the emergency room (ER), and they are responsible for 1 million or 1% of all the ER consults. The annual cost of care, prehospital and within the ER, to treat ES is estimated

to be 1 billion dollars. Clinical guidelines attempt to direct and guarantee the assertiveness of medical attention to improve the diagnosis and treatment of this disease¹⁻³.

Recommendations for scientific research were based on the selective location of keywords in PubMed-MEDLINE, The Cochrane Library, and other clinical practice guidelines, as well as recommendations by other scientific societies.

Correspondence:

Elvira Castro Martínez

E-mail: elviracastrom@yahoo.com.mx

Date of reception: 01-02-2019

Date of acceptance: 28-02-2019

DOI: 10.24875/RMN.M19000025

Disponible en internet: 12-04-2019

Rev Mex Neuroci. 2018;20(2):69-75

www.revmexneurociencia.com

1665-5044/© 2019. Academia Mexicana de Neurología A.C. Published by Permanyer México. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Questions addressed within this guideline

1. What clinical data must be purposefully investigated during the anamnesis of a patient with ES in the ER?
2. Which laboratory tests must be performed in the ER on a patient with a first ES?
3. Under which conditions must electroencephalography be performed on a patient with ES in the ER?
4. Under which conditions must a neuroimaging test (CT or MRI) be performed on a patient with ES in the ER?
5. Under which conditions must anti-epileptic drugs be initiated on patients with a first ES in the ER?

ES: epileptic seizures; ER: emergency room; CT: computerized tomography; MRI: magnetic resonance imaging.

What clinical data must be purposefully investigated during the anamnesis of a patient with ES in the ER?

Recommendation (Figures 1-3)	Level of recommendation
<p>Identification of all possible triggering factors for the seizure:</p> <ul style="list-style-type: none"> – Fever, concomitant systemic infection or disease – History of previous neurological disease – Trauma – Ingestion of drugs or toxic substances – Recent vaccinations – Family history of ES 	<p>D NICE, 2012⁴ R- Solari F, 2011⁵</p>
<p>Medical personnel performing an initial examination on a patient in the ER must identify and interrogate:</p> <ul style="list-style-type: none"> – Alteration of the level of consciousness – Type and topography of altered motor activity – Sensory symptoms – Autonomic symptoms – Cognitive symptoms – Behavior during the preictal, ictal, and postictal periods 	<p>D NICE, 2012⁴ R- SAdE, 2015⁶</p>
<p>Additional information must be gathered for the diagnosis, including:</p> <ul style="list-style-type: none"> – Perinatal pathologies – Characteristics of psychomotor development timeline – Learning disabilities – Other neurological or psychiatric diseases 	<p>3 NICE, 2012⁴ R- SAdE, 2015⁶</p>
<p>A directed neurological examination is recommended in search of signs and symptoms of:</p> <ul style="list-style-type: none"> – Intracranial hypertension – Signs of meningeal irritation – Focal neurological deficit – Neurological emergency 	<p>D NICE, 2012⁴ R- SAdE, 2015⁶</p>

ES: epileptic seizures; ER: emergency room.

Which laboratory tests must be performed in the ER on a patient with a first ES?

Recommendation	Level of recommendation
<p>Laboratory tests on a patient that has complete recovery after a first ES are not considered necessary, unless there are specific findings that justify them</p>	<p>3 NICE, 2012⁴ R-SAdE, 2015⁶</p>
<p>Laboratory tests recommended for patients with ES that present with dehydration due to vomiting or diarrhea and demonstrate a progressive or persistent deterioration of consciousness are:</p> <ul style="list-style-type: none"> – Complete blood count – Blood glucose, sodium, and electrolytes – Serum pregnancy (qualitative hCG) test 	<p>D NICE, 2012⁴ R-SAdE, 2015⁶</p>

Figure 1. Unprovoked Epileptic Seizure.

LP is recommended for any patient with a first ES and suspicion of CNS infection, subarachnoid hemorrhage determined by a non-diagnostic CT image, suspicion of HIV infection, or children under 6 months of age	D NICE, 2012 ⁴ R- SAdE, 2015 ⁶
A LP is not indicated on patients who show complete recovery of their neurological basal state after a first ES	R- SAdE, 2015 ⁶
To establish a differential diagnosis of non-epileptic paroxysmal events, individualization of cases is recommended when considering complementary tests	D NICE, 2012 ⁴ R-SAdE, 2015 ⁶
If substance abuse of psychoactive drugs or exposure to toxic substances is suspected, a toxicology screen is recommended	D NICE, 2012 ⁴ R- SAdE, 2015 ⁶
Complementary diagnostic tests after a new ES on patients with a known history of epilepsy are not necessary, unless there is suspicion of lack of pharmaceutical efficacy, intoxication, adherence to treatment, or change in seizure pattern	R-SAdE, 2015 ⁶

LP: lumbar puncture; ES: epileptic seizures; CNS: central nervous system; CT: computerized tomography.

Figure 2. Acute symptomatic epileptic seizure (provoked).

Under which conditions must electroencephalography be performed on a patient with ES in the ER?

Recommendation	Level of recommendation
There is no evidence that an emergency EEG has any implication on immediate therapy; therefore, there are few indications such as: <ul style="list-style-type: none"> – Suspicion of subtle or non-convulsive status epilepticus – Comatose state of unknown origin – Suspicion of herpetic encephalitis 	IV SAdE, 2015 ⁶
A routine EEG in the ER is not recommended for previously healthy patients (pediatric or adult) that present a first ES and have returned to their basal state	R-SAdE, 2015 ⁶
An EEG is a useful diagnostic test in stabilized patients in the ER to: <ul style="list-style-type: none"> – Support the diagnosis of epilepsy – Determine the type of epilepsy and epileptic syndrome, in concordance with clinical findings and ictal activity – Asses the risk of seizure recurrence – Support therapeutic decisions 	2+ SIGN, 2015 ⁷ 3 NICE, 2012 ⁴ R- SAdE, 2015 ⁶

EEG: electroencephalography; ES: epileptic seizures, ER: emergency room.

Figure 3. Epileptic seizure on patients previously diagnosed with epilepsy.

Under which conditions must a neuroimaging test (CT or MRI) be performed on a patient with ES in the ER?

Recommendation	Level of recommendation
A cranial CT must be performed on all emergency cases; however, it does not substitute a programmed cranial MRI	B SAdE, 2015 ⁶
An emergency cranial CT may be considered for previously healthy patients (pediatric or adult) if a programmed MRI cannot be performed within the following 3 days	R-SAdE, 2015 ⁶

Neuroimaging tests (cranial CT or MRI, the latter being the first choice) are recommended for patients who present a first ES under the following circumstances: <ul style="list-style-type: none"> – Focal seizures – Focal neurological deficit – Persistent alteration in the level of consciousness – Recent traumatic brain injury – Cancer – Suspicion of HIV infection/immunosuppression – Hemorrhagic diathesis – Children under 2 years of age – Persistent cephalgia (headache) – Meningeal signs – Signs of intracranial hypertension 	B SAde, 2015 ⁶ C NICE, 2012 ⁴ R-Ghofrani M, 2013 ⁸ R-Aprahamian N, 2014 ⁹ R-Michoulas AS, 2011 ¹⁰
Neuroimaging tests are not useful under the following circumstances: <ul style="list-style-type: none"> – Febrile seizures – Focal impaired awareness seizures – ES with a proven metabolic origin 	R-SAde, 2015 ⁶

CT: computerized tomography; MRI: magnetic resonance imaging.

Under which conditions must anti-epileptic drugs be initiated on patients with a first ES in the ER?

Recommendation	Level de recommendation
Before initiating treatment with AED on a patient with a first ES, it is recommended to consider: <ul style="list-style-type: none"> – The probability of recurrence – The drugs efficacy and toxicity 	C NICE, 2012 ⁴ R-Bergey G, 2016 ¹¹
Treatment with AED after a first ES must be individualized and assessed by a qualified professional	A NICE, 2012 ⁴ R- Leone MA, 2016 ¹² C NICE, 2012 ⁴ R-Bergey G, 2016 ¹¹
It is recommended that AED treatment be initiated on patients with one or more of the following risk factors: <ul style="list-style-type: none"> – History of previous brain injury with remote symptomatic seizures – Focal ES – ES during sleep – Family history of ES – Abnormal neurological examination – Psychomotor delay – Status epilepticus – Abnormal EEG – Neuroimaging abnormalities 	C NICE, 2012 ⁴ R-Michoulas AS, 2011 ¹⁰ R-Bergey G, 2016 ¹¹ R-Ghofrani M, 2013 ⁸ R-Krumholz A, 2015 ¹³
During the initiation of AED treatment, it is recommended to consider the risk of seizure aggravation: <ul style="list-style-type: none"> – Phenytoin aggravates absence and myoclonic seizures – Carbamazepine and oxcarbamazepine exacerbate absence, myoclonic, and atonic seizures 	D NICE, 2012 ⁴ R-SAde, 2015 ⁶

AED: anti-epileptic drug; EEG: electroencephalography.

This article and its authors do not present any conflict of interest or source of financing.

Algorithm: Management of ES in the ER

References

1. Cerdá JM, Argani MT, Llerda JA, et al. Guía oficial de la sociedad española de neurología de práctica clínica en epilepsia. *Neurología*. 2016;31:121-9.
2. Pallin DJ, Goldstein JN, Moussally JS, et al. Seizure visits in US emergency departments: epidemiology and potential disparities in care. *Int J Emerg Med*. 2008;1:97-105.
3. Martindale JL, Goldstein JN, Pallin DJ. Emergency department seizure epidemiology. *Emerg Med Clin North Am*. 2011;29:15-27.
4. National Clinical Guideline Centre (UK). The Epilepsies: the Diagnosis and Management of the Epilepsies in Adults and Children in Primary and Secondary Care: pharmacological Update of Clinical Guideline 20. London: Royal College of Physicians (UK) (NICE Clinical Guidelines, No. 137); 2012.
5. Solari BF. Crisis epilépticas en la población infantil. *Rev Med Clin Condes*. 2011;22:647-54.
6. Sociedad Andaluza de Epilepsia. Guía Andaluza de Epilepsia. Barcelona: Sociedad Andaluza de Epilepsia; 2015. p. 484.
7. Scottish Intercollegiate Guidelines Network (SIGN). Diagnosis and Management of Epilepsy in Adults. Edinburgh: Scottish Intercollegiate Guidelines Network; 2015.
8. Ghofrani M. Approach to the first unprovoked seizure-PART I. *Iran J Child Neurol*. 2013;7:1-5.
9. Aprahamian N, Harper MB, Prabhu SP, et al. Pediatric first time non-febrile seizure with focal manifestations: is emergent imaging indicated? *Seizure*. 2014;23:740-5.
10. Michoulas A, Farrell K, Connolly M. Approach to a child with a first afebrile seizure. *BCM J*. 2011;53:274-7.
11. Bergey GK. Management of a first seizure. *Continuum (Minneapolis)*. 2016;22:38-50.
12. Leone MA, Giussani G, Nolan SJ, Marson AG, Beghi E. Immediate antiepileptic drug treatment, versus placebo, deferred, or no treatment for first unprovoked seizure. *Cochrane Database Syst Rev*. 2016;5:CD007144.
13. Krumholz A, Wiebe S, Gronseth GS, et al. Evidence-based guideline: management of an unprovoked first seizure in adults: report of the guideline development subcommittee of the American academy of neurology and the American epilepsy society. *Epilepsy Curr*. 2015;15:144-52.